For Research Use Only

SP3 Recombinant antibody

Catalog Number:85050-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

85050-1-RR BC126414 GeneID (NCBI):

Nanodrop: **UNIPROT ID:** Source Q02447 Rabbit Full Name:

100ul, Concentration: 1000 ug/ml by 6670

Isotype: Sp3 transcription factor

IgG Calculated MW: Immunogen Catalog Number: 781 aa, 82 kDa AG25082 Observed MW: 70 kDa, 115 kDa **Purification Method:** Protein A purification

CloneNo.:

242556A2

Recommended Dilutions: WB 1:2000-1:10000 IF/ICC 1:250-1:1000

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA WB: HeLa cells, K-562 cells, PC-3 cells, HCT 116 cells, Species Specificity:

COLO 320 cells, C2C12 cells, C6 cells

IF/ICC: HepG2 cells,

Background Information

SP3, also named as transcription factor Sp3, is a 781 amino acid protein, which contains 3 C2H2-type zinc fingers and belongs to the Sp1 C2H2-type zinc-finger protein family. SP3 localizes to the nuclear periphery and in nuclear dots when sumoylated. Some localization in PML nuclear bodies. SP3 is acetylated by histone acetyltransferase p300, deacetylated by HDACs and sumoylated on all isoforms. Sumoylated on 2 sites in longer isoforms with Lys-551 being the major site. Sumoylation at this site promotes nuclear localization to the nuclear periphery, nuclear dots and PML nuclear bodies. Sumoylation on Lys-551 represses the transactivation activity, except for the largest isoform, L-Sp3, which has little effect on transactivation. Alternate sumoylation and acetylation at Lys-551 also control transcriptional activity. There exits tow modified isoform and molecular weight of those are 70 kda and 115

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

human, mouse, rat

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85050-1-RR (SP3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SP3 antibody (85050-1-RR, Clone: 242556A2) at dilution of 1:500 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).