

For Research Use Only

Caspase 10 Recombinant antibody, PBS Only

Catalog Number: 85196-3-PBS



Basic Information

Catalog Number: 85196-3-PBS	GenBank Accession Number: BC042844	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 843	CloneNo.: 242353H6
Source: Rabbit	UNIPROT ID: Q92851	
Isotype: IgG	Full Name: caspase 10, apoptosis-related cysteine peptidase	
	Calculated MW: 59 kDa	
	Observed MW: 59 kDa	

Applications

Tested Applications:
WB, FC (Intra), Indirect ELISA

Species Specificity:
human

Background Information

CASP10(Caspase-10) is also named as MCH4 and belongs to the peptidase C14A family. It is recruited to the CD99 and p55 tumor necrosis factor receptor signaling complexes and promoting CD95 (APO1/Fas) apoptosis. It may also play a critical role in PTK7-knockdown-induced apoptosis, downstream of mitochondria(PMID:21103379). This protein can be Cleavage by granzyme B and autocatalytic activity generate two active subunits. Defects in CASP10 are the cause of autoimmune lymphoproliferative syndrome type 2A (ALPS2A)(PMID:16446975), familial non-Hodgkin lymphoma (NHL)(PMID:12010812) and gastric cancer (GASC)(PMID:11973654). The full length has a propeptide with 219 amino acid and 6 isoforms produced by alternative splicing.

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85196-3-RR (Caspase 10 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85196-3-PBS in a different storage buffer formulation.



Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 85196-3-RR (Caspase 10 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85196-3-PBS in a different storage buffer formulation.



1×10^6 HepG2 cells were intracellularly stained with 0.25 ug Caspase 10 Recombinant Antibody (85196-3-RR, Clone:242353H6) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 85196-3-PBS in a

