For Research Use Only

Mouse CD8a Recombinant antibody, **PBS Only**

Catalog Number:85977-4-PBS



Purification Method:

CloneNo.:

242879E1

Protein A purification

Basic Information

Catalog Number: 85977-4-PBS

GenBank Accession Number:

NM_001081110.2

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

UNIPROT ID:

Nanodrop:

P01731 Full Name:

Isotype: IgG

CD8 antigen, alpha chain

Immunogen Catalog Number:

Calculated MW: 27 kDa

EG3265

Rabbit

Size:

Observed MW: 30-35 kDa

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

mouse

Background Information

 ${\sf CD8}\ is\ a\ transmembrane\ glycoprotein\ that\ is\ predominantly\ expressed\ on\ the\ surface\ of\ cytotoxic\ T\ cells,\ and\ can$ also be found on natural killer cells, cortical thymocytes, and dendritic cells. CD8 is composed of two disulfidelinked chains and can be present as a homodimer of CD8α or as a heterodimer of CD8α and CD8β (PMID: 3264320; 8253791). The majority of class I-restricted T cells express mostly the CD8a β heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) $\gamma \delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).

Storage

Storage:


Store at -80°C.

Storage Buffer:

PBS only, pH7.3

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85977-4-RR (CD8A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85977-4-PBS in a different storage buffer formulation.