## For Research Use Only

## Anti-Human TNFR2/CD120b Rabbit Recombinant Antibody, PBS Only

Catalog Number:98010-2-PBS





Basic Information	Catalog Number: 98010-2-PBS	GenBank Accession Number: BC052977	Purification Method: Protein A purification
	Size: 1mg , 2 mg/ml	GeneID (NCBI): 7133	CloneNo.: 230328B6
	Source: Rabbit	UNIPROT ID: P20333	23032000
	Isotype: IgG	Full Name: tumor necrosis factor receptor superfamily, member 1B	
		Calculated MW: 48 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	Tumor necrosis factor-alpha (TNFA/TNFSF2) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis (PMID: 16407280). TNFA signals through two distinc cell surface receptors, TNFR1 (TNFRSF1A, CD120a, p55) and TNFR2 (TNFRSF1B, CD120b, p75). TNFR1 is widely expressed, whereas TNFR2 exhibits more restricted expression, being found on CD4 and CD8 T lymphocytes, endothelial cells, microglia, oligodendrocytes, neuron subtypes, cardiac myocytes, thymocytes and human mesenchymal stem cells (PMID: 20489699; 22374304). In contrast to TNFR1, TNFR2 does not have a death domain. TNFR2 only signals for antiapoptotic reactions. However, recent evidence indicates that TNFR2 also signals to induce TRAF2 degradation (PMID: 22374304). Various defects in the TNFR2 pathway, due to polymorphisms in the TNFR2 gene, upregulated expression of TNFR2 and TNFR2 shedding, have been implicated in the pathology of several autoimmune disorders (PMID: 20489699).		
Storage	Storage: Store at -80°C. Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human TNFR2/CD120b Rabbit Recombinant Antibody (98010-2-RR, Clone:23032886) and PE-conjugated Goat Anti-Rabbit IgG, or 0.25 ug Isotype Control. Cells were not fixed. This data was developed using the same antibody clone with 98010-2-PBS in a different storage buffer formulation.

Biolayer interferometry (BLL) kinetic assays of 98010-2-RR against Human TNFR2/CD120b were performed. The affinity constant is below 1 pM.