For Research Use Only

Anti-Human TIGIT Rabbit Recombinant Antibody

Catalog Number: 98023-3-RR



Basic Information

Catalog Number:

98023-3-RR

Size:

100ug, 1 mg/ml Source:

Rabbit Isotype:

type:

T cell immunoreceptor with Ig and ITIM domains

Calculated MW: 26 kDa

NM_173799.4

GeneID (NCBI):

UNIPROT ID:

Q495A1-1 Full Name:

201633

GenBank Accession Number:

Purification Method: Protein A purfication

CloneNo.: 240423A7

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

TIGIT (T-cell immunoreceptor with Ig and ITIM domains), also known as VSIG9 or VSTM3, is an immune receptor expressed on T cells, including Treg and memory subsets, as well as on NK cells (PMID: 19011627). It contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT binds to poliovirus receptor (PVR, also called CD155) with high affinity, and also to PVRL2 (CD112) with lower affinity. The interaction of TIGIT with PVR on dendritic cells increases the secretion of IL-10 and decreases the secretion of proinflammatory cytokine and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells. TIGIT can inhibit NK cytotoxicity directly through its ITIM (PMID: 19815499).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human TIGIT Rabbit Recombinant Antibody (98023-3-RR, Clone: 240423A7) or Isotype Control, and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with APC Anti-Human CD45RO. Cells were not fixed. Lymphocytes were gated.



Biolayer interferometry (BLI) kinetic assays of 98023-3-RR against Human TIGIT were performed. The affinity constant is 4.00 nM.