For Research Use Only

## Anti-Human Neprilysin/CD10 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98116-1-PBS



**Purification Method:** 

Protein A purfication

CloneNo.:

241322E1

**Basic Information** 

Catalog Number:

98116-1-PBS

Size:

1mg, 2 mg/ml Source:

Rabbit Isotype:

GenBank Accession Number:

BC101658 GeneID (NCBI):

4311

**UNIPROT ID:** P08473 Full Name:

membrane metallo-endopeptidase

Calculated MW: 750 aa, 86 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

human


**Background Information** 

CD10, also known as neprilysin, membrane metallo-endopeptidase (MME), neutral endopeptidase (NEP), or common acute lymphoblastic leukemia antigen (CALLA), is a 100-kDa type II transmembrane glycoprotein belonging to peptidase M13 family (PMID: 7760013; 8102558). Among hematopoietic cells, CD10 is expressed on granulocytes, B cell precursors, mature germinal center B cells, a subset of immature thymocytes (PMID: 13679451). CD10 is also expressed on a variety of nonhematopoietic cell types, including bronchial epithelial cells, cultured fibroblasts, bone marrow stromal cells, renal proximal tubular epithelial cells, breast myoepithelium, biliary canaliculi (PMID: 8102558). CD10 is a cell surface peptidase that cleaves peptide bonds on the amino side of hydrophobic amino acids and inactivates a variety of physiologically active peptides. Loss or decreases in CD10 expression have been reported in a variety of malignancies (PMID: 16054017).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Neprilysin/CD10 Rabbit Recombinant Antibody (98116-1-RR, Clone: 241322E1) or Isotype Control and PE-conjugated Goat Anti-Rabbit IgG. Cells were not fixed. This data was developed using the same antibody clone with 98116-1-PBS in a different storage buffer formulation.



1x10^6 human peripheral blood leukocytes were surface stained with 0.25 ug Anti-Human Neprilysin/CD10 Rabbit Recombinant Antibody (98116-1-RR, Clone:241322E1)(red) or Isotype Control (blue) and PE-conjugated Goat Anti-Rabbit IgG. Cells were not fixed. This data was developed using the same antibody clone with 98116-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 98116-1-RR against Human CD10 were performed. The affinity constant is 9.9 nM.