For Research Use Only

## Anti-Human Integrin alpha-5/CD49e Rabbit Recombinant Antibody, PBS Only

Catalog Number:98171-1-PBS



**Purification Method:** 

Protein A purfication

CloneNo.:

241685D4

**Basic Information** 

Catalog Number:

98171-1-PBS

Size: 1mg, 2 mg/ml

Source: Rabbit Isotype:

GenBank Accession Number:

BC008786 GeneID (NCBI):

3678

**UNIPROT ID:** P08648

Full Name:

integrin, alpha 5 (fibronectin receptor,

alpha polypeptide) Calculated MW: 115 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:


## **Background Information**

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated alpha and beta subunits. Integrin alpha-5 (ITGA5, also known as CD49e), belongs to the integrin alpha chain family, primarily binds to Integrin beta 1 (ITGB1) to form an  $\alpha$ 5 $\beta$ 1 heterodimer (PMID: 33711961). Integrin  $\alpha$ 5 $\beta$ 1 is a specific receptor of fibronectin through its arginine-glycine-aspartic acid (RGD) binding site (PMID: 19608542). Integrin α5β1 plays roles in cell adhesion, migration and matrix formation, and has emerged as an essential mediator in many human carcinomas (PMID: 33408483).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human Integrin alpha-5/CD49e Rabbit Recombinant Antibody (98171-1-RR, Clone: 241685D4) or 0.25 ug Isotype Control, and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with APC Anti-Human CD3 (OKT3). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98171-1-PBS in a different storage



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human Integrin alpha-5/CD49e Rabbit Recombinant Antibody (98171-1-RR, Clone: 241685D4) (red) or 0.25 ug Isotype Control (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98171-1-PBS in a different storage buffer formulation.