For Research Use Only

Anti-Human IL-5 Rabbit Recombinant Antibody, PBS Only Catalog Number:98178-1-PBS





| ize: mg , 2 mg/ml ource: abbit | GeneID (NCBI): 3567 UNIPROT ID: | CloneNo.: 240915D12 |
|---|--|---|
| abbit | | |
| cotupo: | P05113 | |
| sotype: gG | Full Name: interleukin 5 (colony-stimulating factor, eosinophil) | |
| | Calculated MW: 134 aa, 15 kDa | |
| | | |
| | | |
| roduced mostly by CD4+ T cell arget cells to activate downstre ole in both innate and adaptive naturation, and survival. Additi ntibody-secreting cells. Elevat | s, eosinophils, and basophils, and signal eam JAK/STAT, Btk, and Ras/ERk pathwa e immune responses. It is a strong promo onally, IL-5 is also capable of stimulatir ed IL-5 levels are strongly correlated wit | s through its specific receptor (IL-5R) on ys. Due to its pleiotropic effects, IL-5 plays a ter of eosinophil differentiation, migration, g activated B cells to differentiate into th eosinophilia, which is a hallmark feature |
| tore at -80°C. torage Buffer: | | |
| | roduced mostly by CD4+ T cell arget cells to activate downstre ole in both innate and adaptive naturation, and survival. Additi ntibody-secreting cells. Elevat | Calculated MW: 134 aa, 15 kDa ested Applications: C (Intra) pecies Specificity: numan L-5 is a 4-0-helix, pleiotropic cytokine that regulates the activities of B d roduced mostly by CD4+ T cells, eosinophils, and basophils, and signal arget cells to activate downstream JAK/STAT, Btk, and Ras/ERk pathway ole in both innate and adaptive immune responses. It is a strong promo- naturation, and survival. Additionally, IL-5 is also capable of stimulatin ntibody-secreting cells. Elevated IL-5 levels are strongly correlated wit of allergy, asthma, and hypersensitivity. (PMID: 21986312, 22174445, 19 storage: itor at -80°C. |

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





K_D=6.89×10⁻¹⁰ K_{off}=1.93×10⁻⁴ K_{on}=2.80×10⁵

0.3

1x10^{^6} untreated or PMA, Ionomycin and Brefeldin A treated human PBMCs were intracellularly stained with 0.25 ug Anti-Human IL-5 Rabbit Recombinant Antibody (98178-1-RR, Clone: 240915D12) and PEconjugated Goat Anti-Rabbit IgG. Cells were then stained with Coralite® Plus 647 Anti-Human CD4. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone

Biolayer interferometry (BLL) kinetic assays of 98178-1-RR against Human IL-5 were performed. The affinity constant is 0.689 nM.

50 nM 25 nM 12.5 nM

6.25 nM 3.125 nM 1.56 nM

0.78 nM

400