For Research Use Only

Anti-Mouse Ki-67 Rabbit Recombinant Antibody

Catalog Number:98224-1-RR





Basic Information	Catalog Number: 98224-1-RR	GenBank Accession Number: NM_001081117.2	Purification Method: Protein A purfication
	Size: 100ug , 1000 µg/ml	GenelD (NCBI): 17345	CloneNo.: 241768E2
	Source: Rabbit	UNIPROT ID: E9PVX6-1	
	Isotype: IgG	Full Name: antigen identified by monoclonal antibody Ki 67	
		Calculated MW: 351kDa	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: mouse		
Background Information	The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki-67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.		
Storage	Storage: Store at 2 - 8°C. Stable for one year a Storage Buffer: PBS with 0.09% sodium azide, pH 7.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 untreated or Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody (98224-1-RR, Clone: 241768E2) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with Coralite® Plus 647 Anti-Mouse CD3. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

1x10^6 Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody (98224-1-RR, Clone: 241768E2) (red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). 1x10^6 untreated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug Anti-Mouse Ki-67 Rabbit Recombinant Antibody