## For Research Use Only

## Anti-Human LILRB3/CD85a Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98237-1-PBS



**Purification Method:** 

Protein A purfication

CloneNo.:

241894C10

**Basic Information** 

Catalog Number:

98237-1-PBS

Size: 1mg, 2 mg/ml

Source: Rabbit

Isotype:

GenBank Accession Number:

BC112198 GeneID (NCBI):

11025

**UNIPROT ID:** 075022 Full Name:

leukocyte immunoglobulin-like

receptor, subfamily B (with TM and

ITIM domains), member 3

Calculated MW: 631 aa, 69 kDa

**Applications** 

**Tested Applications:** 

Species Specificity:

human


## **Background Information**

The leukocyte immunoglobulin-like receptors (LIRs, also known as ILTs, CD85, and LILRs) comprise a family of related immunoregulatory receptors encoded within the leukocyte receptor cluster (LRC) at chromosomal region 19q13.4 (PMID: 11491530). LIRs are transmembrane proteins containing either two or four extracellular immunoglobulin domains, and have diverse functions, including the regulation of inflammation, immune tolerance, cell differentiation and nervous system plasticity (PMID: 16406677; 26040207). LILRB3, also known as ILT-5 or CD85a, belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). LILRB3 is found on the surface of a variety of cell types including monocytes/macrophages, granulocytes, NK cells and some T cells. It binds to MHC class I molecules and transduces a negative signal that inhibits stimulation of an immune response. LILRB3 has been reported as a myeloid cell checkpoint that elicits profound immunomodulation (PMID: 32870822).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human LILRB3/CD85a Rabbit Recombinant Antibody (98237-1-RR, Clone: 241894C10) or 0.25 ug Rabbit 1gG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9), and PE-Conjugated Goat Anti-Rabbit 1gG(H+L). Cells were co-stained with Coralite® Plus 647 Anti-Human CD14. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human LILRB3/CD85a Rabbit Recombinant Antibody (98237-1-RR, Clone: 241894C10)(red) or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L).Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 98237-1-PBS in a