

For Research Use Only

Anti-Human CXCL10/IP-10 Rabbit Recombinant Antibody, PBS Only

Catalog Number: 98265-1-PBS



Basic Information

Catalog Number: 98265-1-PBS	GenBank Accession Number: BC010954	Purification Method: Protein A purification
Size: 1mg, 2 mg/ml	GeneID (NCBI): 3627	CloneNo.: 242013E12
Source: Rabbit	UNIPROT ID: P02778	
Isotype: IgG	Full Name: chemokine (C-X-C motif) ligand 10	
	Calculated MW: 11 kDa	

Applications

Tested Applications:
FC (Intra)

Species Specificity:
human

Background Information

CXCL10 (also known as IP-10) is a member of the CXC chemokine family which binds to the CXCR3 receptor to exert its biological effects. CXCL10 is a 12-kDa protein and constitutes two internal disulfide cross bridges. The predicted signal peptidase cleavage generates a 10-kDa secreted polypeptide with four conserved cysteine residues in the N-terminal. The CXCL10 gene localizes on chromosome 4 at band q21, a locus associated with an acute monocytic/B-lymphocyte lineage leukemia exhibiting translocation of t(4; 11)(q21; q23). CXCL10 mediates leukocyte trafficking, adaptive immunity, inflammation, haematopoiesis and angiogenesis. Under proinflammatory conditions CXCL10 is secreted from a variety of cells, such as leukocytes, activated neutrophils, eosinophils, monocytes, epithelial cells, endothelial cells, stromal cells (fibroblasts) and keratinocytes in response to IFN- γ .

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or IFN gamma and brefeldin A treated human PBMCs were intracellularly stained with 0.25 ug Anti-Human CXCL10/IP-10 Rabbit Recombinant Antibody (98265-1-RR, Clone: 242013E12) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 98265-1-PBS in a different storage buffer formulation.