

For Research Use Only

Anti-Human Nectin-2/CD112 Rabbit Recombinant Antibody

Catalog Number: 98283-1-RR



Basic Information

Catalog Number: 98283-1-RR	GenBank Accession Number: NM_001042724.2	Purification Method: Protein A purification
Size: 100ug , 1000 µg/ml	GeneID (NCBI): 5819	CloneNo.: 241725D3
Source: Rabbit	UNIPROT ID: Q92692-2	
Isotype: IgG	Full Name: poliovirus receptor-related 2 (herpesvirus entry mediator B)	
	Calculated MW: 58 kDa	

Applications

Tested Applications:
FC

Species Specificity:
human

Background Information

Nectin 2, also named as PVRL2, CD112, HVEB, PRR2 and PVRR2, is an adhesion molecule widely expressed in cell lines of different lineages, including hematopoietic, neuronal, endothelial and epithelial cells. CD112 belongs to a new family of immunoglobulin-like molecules that includes four members (CD111, CD112, PRR3 and CD155) sharing an ectodomain made of three Ig domains, of V and C types. CD112 is expressed in the myelo-monocytic and megakaryocytic hematopoietic lineages and the function in hematopoiesis is currently unknown. CD112 is an intercellular homophilic adhesion. CD112 localizes specifically at adherens junctions via its cytoplasmic interaction with the scaffold F-actin binding protein afadin. Disruption of the murine CD112 gene leads to infertility of male mice with morphologically aberrant spermatozoa. CD112 mediates entry of some alphaherpesvirus mutants (also named HveB) via its V domain. CD112 is involved in cell to cell spreading of viruses.

Storage


Storage:
Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide, pH 7.3.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ K562 (left) or Daudi cells (right) were surface stained with 0.25 ug Anti-Human Nectin-2/CD112 Rabbit RecAb (98283-1-RR, Clone: 241725D3) (red) or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed.