

For Research Use Only

Anti-Human CD93 Rabbit Recombinant Antibody

Catalog Number: 98337-1-RR



Basic Information

Catalog Number: 98337-1-RR	GenBank Accession Number: NM_012072.3	Purification Method: Protein A purification
Size: 100ug , 1000 µg/ml	GeneID (NCBI): 22918	CloneNo.: 242464E6
Source: Rabbit	UNIPROT ID: Q9NPY3	
Isotype: IgG	Full Name: CD93 molecule	
	Calculated MW: 69 kDa	

Applications

Tested Applications:
FC

Species Specificity:
human

Background Information

CD93, also known as C1qR or C1qR(p), is a single-pass type I transmembrane glycoprotein and a member of the group 14 of the C-type lectin-like domain (CTLD) superfamily, which also includes endosialin/CD248, thrombomodulin, and CLEC14A (PMID: 31287944). CD93 consists of a C-type lectin-like domain, five EGF-like repeats, a mucin-like domain, a transmembrane domain and a cytoplasmic domain (PMID: 28912033). CD93 is mainly expressed on endothelial cells and has been reported to be expressed by neurons and various cells of the hematopoietic system, including monocytes, neutrophils, B cells, natural killer cells, platelets and hematopoietic stem cells (PMID: 37443812). CD93 plays a role in various physiological processes including angiogenesis, inflammation, and cell adhesion (PMID: 31287944).

Storage

Storage:
Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide, pH 7.3.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD93 Rabbit RecAb (98337-1-RR, Clone: 242464E6) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were then stained with APC Anti-Human CD14. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD93 Rabbit RecAb (98337-1-RR, Clone:242464E6)(red) or Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.