

For Research Use Only

# Anti-Mouse IL-4 Rabbit Recombinant Antibody

Catalog Number: 98360-2-RR



## Basic Information

Catalog Number:

98360-2-RR

Size:

100ug, 1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG3750

GenBank Accession Number:

BC027514

GeneID (NCBI):

16189

UNIPROT ID:

P07750

Full Name:

interleukin 4

Purification Method:

Protein A purification

CloneNo.:

242736G8

Recommended Dilutions:

FC (Intra): 0.13 ug per 10<sup>6</sup> cells in 100 µl suspension

## Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Positive Controls:

FC (Intra): PMA, Ionomycin and Brefeldin A treated C57BL/6 Th2-polarized splenocytes,

## Background Information

Interleukin-4 (IL-4), a member of the α-helical cytokine family, is produced by activated CD4<sup>+</sup> T cells, basophils, and mast cells. It promotes the proliferation and differentiation of antigen-presenting cells. IL-4 also plays a pivotal role in antibody isotype switching and stimulates the production of IgE. This cytokine has been applied in the treatment of autoimmune disorders like multiple myeloma, cancer, psoriasis, and arthritis. IL-4 has also been extensively applied to inhibit detrimental effect of Th1. It may promote the growth of epithelial tumors by mediating increased proliferation and survival. (PMID: 24489573;3049907;21663408)

## Storage

Storage:

Store at 2 - 8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> unstimulated or PMA, Ionomycin and Brefeldin A treated C57BL/6 Th2-polarized splenocytes were intracellularly stained with 0.125 ug Anti-Mouse IL-4 Rabbit RecAb (98360-2-RR, Clone: 242736G8) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with APC Anti-Mouse CD4 (GK1.5) (APC-65104, Clone: GK1.5). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.