

For Research Use Only

APC Anti-Human TNFR1/CD120a Rabbit Recombinant Antibody

Catalog Number: APC-98085



Basic Information

Catalog Number: APC-98085	GenBank Accession Number: BC010140	Purification Method: Protein A purification
Size: 100tests , 5 ul/test	GeneID (NCBI): 7132	CloneNo.: 241337H6
Source: Rabbit	UNIPROT ID: P19438	Excitation/Emission maxima wavelengths: 650 nm / 660 nm
Isotype: IgG	Full Name: tumor necrosis factor receptor superfamily, member 1A	
	Calculated MW: 455 aa, 50 kDa	

Applications

Tested Applications:
FC

Species Specificity:
human

Background Information

Tumor necrosis factor (TNF) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis (PMID: 16407280). TNF exists in soluble and membrane-bound forms. TNF signals through two distinct cell surface receptors, TNFR1 (TNFRSF1A, CD120a) and TNFR2 (TNFRSF1B, CD120b). Whereas TNFR1 is widely expressed, expression of TNFR2 is limited to cells of the immune system, endothelial cells, and nerve cells (PMID: 22053109). TNFR1, which contains a death domain (DD) within its intracytoplasmic region, is thought to be the key receptor for TNF signaling (PMID: 16407280). This receptor can activate NF-kappaB, mediate apoptosis, and function as a regulator of inflammation. Antiapoptotic protein BCL2-associated athanogene 4 (BAG4/SODD) and adaptor proteins TRADD and TRAF2 have been shown to interact with this receptor, and thus play regulatory roles in the signal transduction mediated by the receptor.

Storage


Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:
PBS with 0.09% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human peripheral blood leukocytes were surface stained with 5 ul APC Anti-Human TNFR1/CD120a Rabbit RecAb (APC-98085, Clone: 241337H6) or APC Rabbit IgG Isotype Control RecAb (APC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10⁶ human peripheral blood leukocytes were surface stained with 5 ul APC Anti-Human TNFR1/CD120a Rabbit RecAb (APC-98085, Clone: 241337H6) (red) or APC Rabbit IgG Isotype Control RecAb (APC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.