For Research Use Only

APC Anti-Human CD45 Rabbit Recombinant Antibody

Catalog Number: APC-98117





Basic Information	Catalog Number: APC-98117 Size: 100tests , 5 ul/test Source: Rabbit Isotype: IgG	GenBank Accession Number: GeneID (NCBI): 5788 ENSEMBL Gene ID: ENSG00000081237 Full Name: protein tyrosine phosphatase, receptor type, C	Purification Method: Protein A purification CloneNo.: 241670E10 Excitation/Emission maxima wavelengths: 650 nm / 660 nm
Applications	Tested Applications: FC Species Specificity: human		
Background Information	CD45, also known as protein tyrosine phosphatase, receptor type C, is a type I transmembrane protein expressed on the surface of all haematopoietic cells with the exception of erythrocytes and platelets (PMID: 3489673; 28615666). CD45 is a pan-haematopoietic cell marker and has been shown to be essential for T- and B-cell activation and signalling (PMID: 9429890; 16378097).		
Storage	Storage: Store at 2-8°C. Avoid exposure to li Storage Buffer: PBS with 0.1% sodium azide and 0.	ght. Stable for one year after shipment 5% BSA.	t.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 human peripheral blood leukocytes were surface stained with 5 ul APC Anti-Human CD45 Rabbit Recombinant Antibody (APC-98117, Clone: 241670E10) or APC Rabbit IgG Isotype Control Recombinant Antibody (APC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.

1x10^6 human peripheral blood leukocytes were surface stained with 5 ul APC Anti-Human CD45 Rabbit Recombinant Antibody (APC-98117, Clone:241670E10) (red) or APC Rabbit IgG Isotype Control Recombinant Antibody (APC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.