

For Research Use Only

Biotin Plus-conjugated ATP1A2-Specific Polyclonal antibody



Catalog Number: Biotin-55179

Featured Product

Basic Information

Catalog Number:

Biotin-55179

GenBank Accession Number:

NM_000702

Purification Method:

Antigen Affinity Purified

Size:

100ul, Concentration: 1000 ug/ml by 477

GeneID (NCBI):

477

Recommended Dilutions:

IHC 1:50-1:500

Nanodrop;

UNIPROT ID:

P50993

Source:

Rabbit

Full Name:

ATPase, Na⁺/K⁺ transporting, alpha 2 (+) polypeptide

Isotype:

IgG

Calculated MW:

112 kDa

Observed MW:

100 kDa

Applications

Tested Applications:

IHC

Positive Controls:

IHC : mouse heart tissue,

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

ATP1A2, also named as KIAA0778, belongs to the cation transport ATPase (P-type) family and Type IIC subfamily. It is the catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of sodium and potassium ions across the plasma membrane. Defects in ATP1A2 are the cause of familial hemiplegic migraine type 2 (FHM2). Defects in ATP1A2 are a cause of alternating hemiplegia of childhood (AHC). This antibody is specific to ATP1A2.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse heart tissue slide using Biotin-55179 (ATP1A2-Specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).