For Research Use Only

Biotin Anti-Human CD16 (3G8) Mouse IgG2a Recombinant Antibody

Catalog Number:Biotin-65612



Basic Information

Catalog Number: Biotin-65612

BC017865

Purification Method: Protein A purification

GeneID (NCBI):

CloneNo.:

100ug , 500 μ g/ml

3G8

Source: Mouse

ENSEMBL Gene ID: ENSG00000203747

Full Name: Isotype: lgG2a

Fc fragment of IgG, low affinity IIIa,

receptor (CD16a)

GenBank Accession Number:

Calculated MW: 254 aa, 29 kDa

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD16 is a 50-70-kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil $polymorphonuclear\ leukocytes, monocytes\ and\ macrophages.\ CD16\ mediates\ antibody-dependent\ cellular$ $cytotoxicity \ (ADCC) \ and \ other \ antibody-dependent \ responses, such as \ phagocytosis. \ CD16 \ has \ been \ identified \ as \ Fc$ receptors Fcy RIIIa (CD16a) and Fcy RIIIb (CD16b), encoded by two nearly identical genes, FCGR3A and the FCGR3B. Clone 3G8 recognizes both the CD16a and CD16b (PMID: 7592758).


Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide.

Selected Validation Data





1x10^6 human peripheral blood leukocytes were surface stained with 5 ul Biotin Plus Anti-Human CD16 (3G8) Mouse IgG2a Recombinant Antibody (Biotin-65612, Clone: 3G8) and Coralite®488-conjugated streptavidin, or unstained cells. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human peripheral blood leukocytes were surface stained with 5 ul Biotin Plus Anti-Human CD16 (3G8) Mouse IgG2a Recombinant Antibody (Biotin-65612, Clone: 3G8) and Coralite@488-conjugated streptavidin(red), or unstained cells(blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.