For Research Use Only

## CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a Recombinant Antibody



Catalog Number: CL405-65592

**Basic Information** 

Catalog Number: CL405-65592

Size:

100tests, 5 ul/test

Source: Mouse

Isotype: lgG2a

GenBank Accession Number:

BC040261 GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000114013

Full Name: CD86 molecule

Calculated MW: 329 aa, 38 kDa

**Purification Method:** 

Protein A purification

CloneNo.: BU63

Excitation/Emission maxima

wavelengths: 399 nm / 422 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

**Background Information** 

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding  $of CD86\ with\ CD28\ antigen\ is\ a\ costimulatory\ signal\ for\ activation\ of\ the\ T-cell.\ Binding\ of\ CD86\ with\ cytotoxic\ T-cell.$  $lymphocyte-associated\ protein\ 4\ negatively\ regulates\ T-cell\ activation\ and\ diminishes\ the\ immune\ response.$ 

Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide.

## **Selected Validation Data**





1x10^6 human PBMCs were surface stained with APC Anti-Human CD14 Rabbit Recombinant Antibody (APC-98040, Clone: 230332D7), and 5 ul CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL405-6559, Clone: BU63) or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 405 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL405-65592, Clone:BU63) (red) or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.