For Research Use Only

CoraLite® Plus 488-conjugated EIF2A/CDA02 Polyclonal antibody Catalog Number:CL488-11233 Featured Product





Basic Information	Catalog Number: CL488-11233	GenBank Accession Number: BC011885	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	100ul , Concentration: 1000 ug/ml b	y 83939	IF/ICC 1:50-1:500	
	Nanodrop;	UNIPROT ID: Q9BY44 Full Name:	Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
	Source:			
	Rabbit			
	lsotype:	eukaryotic translation initiation factor 2A, 65kDa Calculated MW: 65 kDa Observed MW: 65 kDa		
	IgG			
	Immunogen Catalog Number: AG1753			
		Applications	Tested Applications:	Tested Applications: Positive Controls:
IF/ICC, FC (Intra)	IF/ICC : HepG2 cells,			
Species Specificity:				
human, mouse, rat				
Packaround Informati	FIF2A, also named as CDA02, MSTP0	004 and MSTP089, belongs to the WI) repeat FIF2A family. It functions in the	
Background Information	early steps of protein synthesis of a protein, has a key role in the transla phosphorylation by activating the el residues near the C terminus of hum 83939) and EIF2S1 (Gene ID: 1965) s translation initiation complex. Altho	EIF2A, also named as CDA02, MSTP004 and MSTP089, belongs to the WD repeat EIF2A family. It functions in the early steps of protein synthesis of a small number of specific mRNAs. EIF2A, an alternative initiator tRNA-binding protein, has a key role in the translation of HCV mRNA during HCV infection, in turn promoting eIF2a phosphorylation by activating the eIF2a kinase PKR. This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human EIF2A, recognizes specifically the 65kd EIF2A protein. EIF2A (Gene ID: 83939) and EIF2S1 (Gene ID: 1965) share the EIF2A symbol/alias in common. EIF2S1 is the alpha subunit of the eIF2 translation initiation complex. Although both of these proteins function in binding initiator tRNA to the 40S ribosomal subunit, the EIF2A protein does so in a codon-dependent manner, whereas eIF2 complex requires GTP.		
Storage	Storage Buffer: PBS with 50% Glycerol, 0.05% Procl	Store at -20°C. Avoid exposure to light. Stable for one year after shipment.		

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite® Plus 488 Eif2a/Cda02 Polyclonal Antibody (CL488-11233)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit JgC control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 EIF2A/CDA02 antibody (CL488-11233) at dilution of 1:200.