## For Research Use Only

## CoraLite® Plus 488-conjugated COXIV Polyclonal antibodv

Polyclonal antic Catalog Number:CL488-1124			www.ptglab.com
Basic Information	Catalog Number: CL488-11242	GenBank Accession Number: BC021236	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1640	GeneID (NCBI): 1327	Recommended Dilutions: IF/ICC 1:50-1:500
		UNIPROT ID: P13073	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
		Full Name:	
		:ytochrome c oxidase subunit IV soform 1 Calculated MW: 19.6 kDa	
		Applications	Tested Applications:
IF/ICC, FC (Intra) Species Specificity: human, mouse, rat	IF/ICC : Her		łepG2 cells,
Background Information	COX411, also named as COX4 and COXIV-1, belongs to the cytochrome c oxidase IV family. It is one of the nuclear- coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport. COX411 is a marker for mitochondria. It has two isoforms (isoform 1 and 2). Isoform 1(COX411) is ubiquitously expressed and isoform 2 is highly expressed in lung tissues. COX411 is commonly used as a loading control. This antibody was generated against full length COX411 protein and cross reacts with COX412.		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	n300, 0.5% BSA, pH 7.3.	t.

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


proteinte

Antibodies | ELISA kits | Proteins

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 COXIV antibody (CL488-11242) at dilution of 1:200.



1X10<sup>^</sup>6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human COXIV (CL488-11242) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).