For Research Use Only

CoraLite® Plus 488-conjugated NAC1 Polyclonal antibody

www.ptglab.com

Purification Method:

Catalog Number: CL488-14551

Catalog Number: **Basic Information**

GenBank Accession Number:

CL488-14551 BC055396 Antigen affinity purification GeneID (NCBI): Recommended Dilutions: IF/ICC 1:50-1:500

100ul , Concentration: 1000 $\mu g/ml$ by 112939 Nanodrop: **UNIPROT ID:** Excitation/Emission maxima

Q96RE7 wavelengths: 493 nm / 522 nm Rabbit Full Name:

Isotype: nucleus accumbens associated 1, BEN and BTB (POZ) domain containing IgG

Immunogen Catalog Number: Calculated MW:

AG6096 57 kDa

Observed MW: 60 kDa

Applications

Tested Applications:

Species Specificity: human, mouse

Positive Controls:

IF/ICC: MCF-7 cells,

Background Information

NACC1, also named as BTBD14B and NAC1, belongs to the BTB/POZ family. NACC1 is a transcriptional repressor, which is amplified and overexpressed in various human cancers and plays critical roles in tumor development, progression, and drug resistance (PMID: 31101655). NACC1 has 1 isoform with the molecular mass of 57 kDa. While NAC1 has been shown to repress the transcription of two tumor suppressors, Gadd45-g and Gadd45-g-interacting protein 1, scientists have found that NACC1 also plays an important role in nontranscriptional function, for example, it can act as a bridge for MAVS and TBK1 in RLR-mediated signaling (PMID: 31235549).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Coralite® Plus 488 NAC1 antibody (CL488-14551) at dilution of 1:200, CL594-Phalloidin (red).