For Research Use Only

CoraLite® Plus 488-conjugated CD146/MCAM Polyclonal antibody



Purification Method:

IF/ICC 1:200-1:800

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number:CL488-17564 Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-17564 BC056418 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 4162 Nanodrop: **UNIPROT ID:**

P43121 Rabbit Full Name:

Isotype: melanoma cell adhesion molecule

IgG Calculated MW: Immunogen Catalog Number: 646 aa, 72 kDa AG11762 Observed MW:

113-120 kDa

Applications

Tested Applications:

IF/ICC

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HUVEC cells,

Background Information

CD146, also known as melanoma cell adhesion molecule (MCAM) or MUC18, originally identified as a biomarker of melanoma progression, is a transmembrane glycoprotein of 113-130 kDa, belonging to the immunoglobulin (Ig) superfamily (PMID: 8378324; 25993332). Structurally, it consists of five Ig domains, a transmembrane domain, and a cytoplasmic region. In normal adult tissue, CD146 is primarily expressed by vascular endothelium and smooth muscle. CD146 is a key cell adhesion protein in vascular endothelial cell activity and angiogenesis, and has been used as marker of circulating endothelium cells (CECs) (PMID: 19356677). In addition to the membrane-anchored form of CD146, a soluble form of CD146 (sCD146, 105 kDa) has also been found in human plasma and in the supernatant of cultured human endothelial cells (PMID: 9462829; 19229070; 16374253; 14597988).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using CoraLite® Plus 488 CD146/MCAM antibody (CL488-17564) at dilution of 1:400.