For Research Use Only

CoraLite® Plus 488-conjugated DDX17 Polyclonal antibody

www.ptglab.com

Purification Method:

IF/ICC 1:50-1:500

493 nm / 522 nm

wavelengths:

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-19910

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-19910 BC000595 GeneID (NCBI): 100ul, Concentration: 1000 ug/ml by 10521

Nanodrop: **UNIPROT ID:** Q92841 Rabbit Full Name:

Isotype: DEAD (Asp-Glu-Ala-Asp) box

polypeptide 17 IgG Immunogen Catalog Number: Calculated MW: AG13723 729 aa. 80 kDa Observed MW:

72-80 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)

human, mouse, rat

Positive Controls: IF/ICC: HEK-293 cells, Species Specificity:

Background Information

DDX17, also named as DEAD box protein p72 and DEAD box protein p82, is a 729 amino acid protein, which belongs to the DEAD box helicase family. DDX5/DBP2 subfamily. DDX17 is widely expressed. Levels tend to increase during colon cancer progression, from very low in benign hyperplastic polyps to very high in tubular and villous adenomas. DDX17 as an RNA helicase unwinds RNA and alters RNA structures through ATP binding and hydrolysis. DDX17 is involved in multiple cellular processes, including pre-mRNA splicing, alternative splicing, ribosomal RNA processing and miRNA processing, as well as transcription regulation. It Regulates the alternative splicing of exons exhibiting specific features. identification and characterisation of p72, a novel human nuclear DEAD box protein, which shows a striking homology to p68. a predicted molecular mass of DDX17 is 72 kDa, but endogenous DDX17 migrates aberrantly at 79 kDa on SDS-PAGE (PMID: 8871553).


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using Coralite® Plus 488 DDX17 antibody (CL488-19910) at dilution of 1:200, CL594-Phalloidin (red).



1x10^6 HEK-293 cells were intracellularly stained with 0.8 ug Coralite® Plus 488-conjugated DDX17 Polyclonal antibody (CL488-19910)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-3000) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).