

For Research Use Only

# CoraLite® Plus 488-conjugated Osteopontin Polyclonal antibody

Catalog Number: CL488-22952

Featured Product

1 Publications



## Basic Information

<b>Catalog Number:</b> CL488-22952	<b>GenBank Accession Number:</b> BC007016	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 6696	<b>Excitation/Emission maxima wavelengths:</b> 493 nm / 522 nm
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P10451	
<b>Isotype:</b> IgG	<b>Full Name:</b> secreted phosphoprotein 1	
<b>Immunogen Catalog Number:</b> AG19216	<b>Calculated MW:</b> 314 aa, 35 kDa	
	<b>Observed MW:</b> 70 kDa, 44-66 kDa	

## Applications

**Tested Applications:**  
FC (Intra)

**Cited Applications:**  
IF

**Species Specificity:**  
human, mouse, rat

## Background Information

Osteopontin (OPN), also known as SPP1, is a secreted glycoposphoprotein that belongs to the small integrin-binding ligand N-linked glycoprotein (SIBLING) family. Originally isolated from bone, OPN has been found in kidney, vascular tissues, biological fluids, and various tumor tissues (PMID: 15138464; 16406521). OPN can interact with CD44 and integrins and regulate diverse biological processes. It has a multifaceted role in bone development and remodeling, and is also involved in the inflammatory and immune response, oncogenesis and cancer progression. The very acidic nature of OPN, as well as the presence of variable posttranslational modifications, has led to anomalous migration in SDS-polyacrylamide gels and therefore to reports of different molecular weights for OPN (PMID: 8293561). Depending on the cell and tissue source and/or the SDS-PAGE system, OPN migrates with a molecular weight of 44-80 kDa, as well as at some smaller bands correspond to peptide fragments (PMID: 8195113; 17890765).

## Notable Publications

Author	Pubmed ID	Journal	Application
Weibo Huang	39326787	J Lipid Res	IF

## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human Osteopontin (CL488-22952) (red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000, Clone: ) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).