For Research Use Only

CoraLite® Plus 488-conjugated OGT Monoclonal antibody

Catalog Number:CL488-66823 Featured Product




| Basic Information | Catalog Number: | GenBank Accession Number: | Purification Method: | | |
|------------------------|--|---|---|--------------------------------------|------------------------------------|
| | CL488-66823 | BC014434 | Protein A purification | | |
| | Size: | GeneID (NCBI): | CloneNo.: | | |
| | 100ul , Concentration: 1000 µg/ml by Nanodrop; Source: Mouse Isotype: IgG2a | | 2B2A6 | | |
| | | UNIPROT ID: | Recommended Dilutions: | | |
| | | O15294 Full Name: O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N- acetylglucosamine:polypeptide-N- acetylglucosaminyl transferase) | WB 1:1000-1:4000 Excitation/Emission maxima wavelengths: 493 nm / 522 nm | | |
| | | | | Immunogen Catalog Number: AG28402 | |
| | | | | | Calculated MW: 1046 aa, 117 kDa |
| | Observed MW: 110 kDa | | | | |
| | Applications | Tested Applications: | Positive Controls: WB : HepG2 cells, | | |
| | | WB | | | |
| | | Species Specificity: human, mouse, rat, pig | | | |
| Background Information | O-linked N-acetylglucosamine transferase (OGT) catalyzes the attachment of N-acetylglucosamine (GlcNAc) monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression. O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues. | | | | |
| | | | | | |
| Storage | Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage | | | | |
| | | | | | |

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL488-66823 (OGT antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.