

For Research Use Only

CoraLite® Plus 488-conjugated Hexokinase 2 Monoclonal antibody

Catalog Number:CL488-66974



Basic Information

Catalog Number: CL488-66974	GenBank Accession Number: BC021116	Purification Method: Protein G purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 3099	CloneNo.: 2A11C3
Source: Mouse	UNIPROT ID: P52789	Recommended Dilutions: WB 1:1000-1:8000
Isotype: IgG1	Full Name: hexokinase 2	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Immunogen Catalog Number: AG16895	Calculated MW: 102 kDa	
	Observed MW: 102 kDa	

Applications

Tested Applications: WB, FC (Intra)	Positive Controls: WB : HeLa cells, HepG2 cells
Species Specificity: human, mouse, rat	

Background Information

Hexokinase 2 (HK2), a rate-limiting enzyme in the first step of glycolysis pathway, expresses at high level in cancer cells compared with normal cells. HK2 provides a new target for cancer therapy due to its pivotal role in tumor tumorigenic and metastatic process. HK1 is constitutively expressed in most mammalian adult tissues. HK2, however, although is abundantly expressed in embryonic tissues, is expressed at high levels only in limited number of adult tissues such as adipose, skeletal, and cardiac muscles. (PMID: 29305912, PMID: 28427443, PMID: 23911236)

Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 HepG2 cells were intracellularly stained with 0.4 μ g Coralite® Plus 488 Hexokinase 2 Monoclonal Antibody (CL488-66974, Clone:2A11C3)(red), or 0.4 μ g CoraLite® Plus 488 Mouse IgG1 Isotype Control (MOPC-21) (CL488-65124, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Various lysates were subjected to SDS PAGE followed by western blot with CL488-66974 (Hexokinase 2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.