For Research Use Only

CoraLite® Plus 488-conjugated VDAC1/Porin Recombinant antibody

Catalog Number: CL488-81538




Basic Information	Catalog Number: CL488-81538	GenBank Accession Number: NM_003374	Purification Method: Protein A purification	
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI): 7416	CloneNo.: 2F4	
		UNIPROT ID: P21796 Full Name: voltage-dependent anion channel 1	Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
				Calculated MW: 31 kDa
		Observed MW: 31 kDa		
		Applications	Tested Applications: FC (Intra)	
Species Specificity: human, mouse, rat				
Background Information	VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.			
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C st	n300, 0.5% BSA, pH 7.3.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite® Plus 488 Vdac1/Porin Recombinant Antibody (CL488-81538, Clone:2F4) (red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).