

For Research Use Only

# CoraLite® Plus 488-conjugated GAPDH Recombinant antibody

Catalog Number:CL488-81640-5



## Basic Information

|  |   |  |
|--|---|--|
| <b>Catalog Number:</b><br>CL488-81640-5                        | <b>GenBank Accession Number:</b><br>BC004109                  | <b>Purification Method:</b><br>Protein A purification              |
| <b>Size:</b><br>100ul , Concentration: 1000 µg/ml by Nanodrop; | <b>GeneID (NCBI):</b><br>2597                                 | <b>CloneNo.:</b><br>1H18   |
| <b>Source:</b><br>Rabbit                                       | <b>UNIPROT ID:</b><br>P04406                                  | <b>Recommended Dilutions:</b><br>IF/ICC 1:50-1:500                 |
| <b>Isotype:</b><br>IgG   | <b>Full Name:</b><br>glyceraldehyde-3-phosphate dehydrogenase | <b>Excitation/Emission maximum wavelengths:</b><br>493 nm / 522 nm |
| <b>Immunogen Catalog Number:</b><br>AG0766                     | <b>Calculated MW:</b><br>36 kDa                               |  |
|  | <b>Observed MW:</b><br>36 kDa                                 |  |

## Applications

|  |   |
|--|---|
| <b>Tested Applications:</b><br>IF/ICC            | <b>Positive Controls:</b><br>IF/ICC : HeLa cells, |
| <b>Species Specificity:</b><br>human, mouse, rat |   |

## Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 GAPDH antibody (CL488-81640-5, Clone: 1H18 ) at dilution of 1:200.