For Research Use Only

CoraLite®555-conjugated LBR Polyclonal antibody

Antibodies | ELISA kits | Proteins www.ptglab.com


Catalog Number: CL555-12398

Basic Information	Catalog Number: CL555-12398	GenBank Accession Number: BC020079	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;		Excitation/Emission maxima wavelengths: 557 nm / 570 nm
	Source: Rabbit Isotype: IgG	Full Name: 357 min 7 576 min lamin B receptor Calculated MW: 615 aa, 71 kDa Observed MW: 58 kDa 58 kDa	
			Applications
Species Specificity: human, mouse, rat			
Background Information	Lamin-B receptor (LBR) is an integral membrane protein of the inner nuclear membrane that contains a hydrophilic N-terminal end protruding into the nucleoplasm, eight hydrophobic segments that span the membrane and a short, nucleoplasmic C-terminal tail (PMID: 28858257). LBR anchors the lamina and the heterochromatin to the inner nuclear membrane (PMID: 10828963; 28858257). It is also essential for cholesterol synthesis (PMID: 27336722). Mutations of the LBR gene have been associated with autosomal recessive HEM/Greenberg skeletal dysplasia and Pelger-Huët anomaly and Greenberg skeletal dysplasia (PMID: 12618959; 12490533). The calculated molecular mass of LBR is 71 kDa, which is larger than the apparent molecular mass of 58 kDa, probably due to the aberrant migration of membrane proteins subjected to SDS-PAGE (PMID: 2847165; 2170422).		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin	, , , , , , , , , , , , , , , , , , ,	nt.
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20°C st		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10[^]6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human LBR (CL555-12398) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).