For Research Use Only

CoraLite®555-conjugated VDAC1/Porin Polyclonal antibody

Catalog Number: CL555-55259

Featured Product

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF-P: 1:50-1:500 IF-Fro: 1:50-1:500

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Basic Information

Catalog Number: CL555-55259

Size: GeneID (NCBI): 100ul , Concentration: 1000 ug/ml by 7416

Nanodrop; UNIPROT ID: Source: P21796

Rabbit Full Name:

Isotype: voltage-dependent anion channel 1
IgG Calculated MW:

31 kDa Observed MW: 31 kDa

NM 003374

GenBank Accession Number:

Applications

Tested Applications: IF-P, IF-Fro, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF-P: human liver cancer tissue,
IF-Fro: mouse liver tissue,
FC (Intra): HepG2 cells,

Background Information

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite®555 VDAC1/Porin antibody (CL555-55259) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite®555-conjugated VDAC 1/Porin Polyclonal antibody (CL555-55259) (red), or 0.8 ug CoraLite®555-conjugated Rabbit 1gG control Rabbit PolyAb (CL555-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse liver tissue using CoraLite®555 VDAC1/Porin antibody (CL555-55259) at dilution of 1:200.