For Research Use Only

CoraLite®594-conjugated Vimentin Polyclonal antibody



Catalog Number:CL594-10366

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL594-10366 BC000163 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7431 Nanodrop: ENSEMBL Gene ID: ENSG00000026025 Rabbit **UNIPROT ID:** Isotype:

P08670 IgG Full Name: Immunogen Catalog Number: vimentin AG0489 Calculated MW:

466 aa, 54 kDa Observed MW: 54 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions: IF/ICC 1:250-1:1000

Excitation/Emission maxima wavelengths: 588 nm / 604 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells,

Background Information

Vimentin, also named as VIM, belongs to the intermediate filament family. Vimentin is class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is important for stabilizing the architecture of the cytoplasm. Monocyte-derived macrophages secrete vimentin into the extracellular space in vitro. Secretion of vimentin was enhanced by the proinflammatory cytokine tumor necrosis factor-alpha (TNFA; 191160) and inhibited by the antiinflammatory cytokine IL10 (124092), suggesting that vimentin is involved in the immune response. Vimentin has specialized functions that contribute to specific dynamic cellular processes. As a phosphoprotein, 55-60 kDa of vimentin proteins can be observed due to the different phosphorylation level. Isoforms of vimentin (49 kDa and 60 kDa) had also been reported. (PMID: 8640945, 22728585).

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 Vimentin antibody (CL594-10366) at dilution of 1:500.



1x10^6 mMSCs cells were intracellularly stained with 0.25 ug CoraLite®594-conjugated Vimentin Polyclonal antibody (CL594-10366)(red). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).