## For Research Use Only

## CoraLite®594-conjugated MMP9 (Nterminal) Polyclonal antibody



Catalog Number:CL594-10375

Featured Product


1 Publications


Basic Information	Catalog Number: CL594-10375	GenBank Accession Number: BC006093	Purification Method: Antigen affinity purification	
	Size: 100ul , Concentration: 1000 µg/ml by	GenelD (NCBI):	Recommended Dilutions: IF 1:50-1:500	
	Nanodrop;			
	Source:	Full Name: matrix metallopeptidase 9	Excitation/Emission maxima wavelengths:	
	Rabbit	(gelatinase B, 92kDa gelatinase,	588 nm / 604 nm	
	Isotype:	92kDa type IV collagenase)		
	lgG	Calculated MW:		
	Immunogen Catalog Number: AG0552	707 aa, 78 kDa		
		Observed MW:		
		67 kDa, 92 kDa		
Applications	Tested Applications:	Positive Co	ontrols:	
Applications	FC (Intra), IF	IF : HepG2 cells,		
	Cited Applications:	тг : перод		
	IF			
	Species Specificity:			
	human, mouse, rat			
	Cited Species:			
	human			
Background Information	normal physiological processes, such processes, such as arthritis or metasta cleaved by extracellular proteinases. collagenase) (MMP9, synonyms: GELB that MMP9 is involved in IL-8-induced	as embryonic development, reproc sis. Most MMP's are secreted as ina Matrix metalloproteinase 9 (gelati , CLG4B) degrades collagens type I mobilization hematopoietic proge ated tissue remodeling. The pro-M	nase B, 92kDa gelatinase, 92kDa type IV V and V. Studies in rhesus monkeys sugge enitor cells from bone marrow, and murine MP9 is 92 kDa, and it can be detected a	
Background Information Notable Publications	normal physiological processes, such processes, such as arthritis or metasta cleaved by extracellular proteinases. collagenase) (MMP9, synonyms: GELB that MMP9 is involved in IL-8-induced studies suggest a role in tumor-associ processed form of 68 kDa. This protein	as embryonic development, reproc sis. Most MMP's are secreted as ina Matrix metalloproteinase 9 (gelati , CLG4B) degrades collagens type I mobilization hematopoietic proge ated tissue remodeling. The pro-M	duction, tissue remodeling, and disease ctive proproteins which are activated whe nase B, 92kDa gelatinase, 92kDa type IV V and V. Studies in rhesus monkeys sugge entor cells from bone marrow, and murine MP9 is 92 kDa, and it can be detected a	
	normal physiological processes, such processes, such as arthritis or metasta cleaved by extracellular proteinases. collagenase) (MMP9, synonyms: GELB that MMP9 is involved in IL-8-induced studies suggest a role in tumor-associ processed form of 68 kDa. This protein	as embryonic development, reprod sis. Most MMP's are secreted as ina Matrix metalloproteinase 9 (gelati , CLG4B) degrades collagens type I I mobilization hematopoietic proge lated tissue remodeling. The pro-M n can exist as a dimer of 180 kDa (F	duction, tissue remodeling, and disease ctive proproteins which are activated whe nase B, 92kDa gelatinase, 92kDa type IV V and V. Studies in rhesus monkeys sugge enitor cells from bone marrow, and murine MP9 is 92 kDa, and it can be detected a MID:7492685).	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human MMP9 (Nterminal) (CL594-10375) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 MMP9 (Nterminal) antibody (CL594-10375) at dilution of 1:200.