For Research Use Only

## CoraLite®594-conjugated GAPDH Monoclonal antibody

Catalog Number:CL594-60004

Featured Product







Basic Information	Catalog Number: CL594-60004	GenBank Accession Number: BC004109	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	100ul , Concentration: 1000 ug/ml by	2597	1E6D9	
	Nanodrop;	UNIPROT ID:	Recommended Dilutions:	
	Source: Mouse	P04406	WB 1:5000-1:50000	
	Isotype: IgG2b	Full Name: glyceraldehyde-3-phosphate dehydrogenase	Excitation/Emission maxima wavelengths: 588 nm / 604 nm	
	Immunogen Catalog Number: AG0766	Calculated MW: 36 kDa		
		Observed MW: 36 kDa		
Applications	Tested Applications: WB, FC (Intra)	Positive Controls:		
	Cited Applications: WB	WB : HeLa cells, HEK-293 cells, Jurkat cells		
	Species Specificity: human, mouse, rat, zebrafish, yeast, plant			
	Cited Species: human, mouse, rat			
Background Information	Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3- phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform o spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.			
	also loading control in electrophoresi membrane, but can occasionally tran such as S-nitrosylation. This antibody kDa GAPDH protein in most cells/tiss spliced product of GAPDH (PMID: 238)	is and Western blotting. GAPDH is slocate to the nucleus after the ac r is raised against full length GAP ues. In addition, a band below 36 85286, 23877755, 19368702). Plea	normally expressed in cellular cytoplasm o Idition of post-translational modifications DH of human origin. It can recognize the 36 kDa can always be detected as the isoform of ase note that some physiological factors, suc	
Notable Publications	also loading control in electrophoresi membrane, but can occasionally tran such as S-nitrosylation. This antibody kDa GAPDH protein in most cells/tiss spliced product of GAPDH (PMID: 238 as hypoxia and diabetes, increase GA	is and Western blotting. GAPDH is slocate to the nucleus after the ac r is raised against full length GAP ues. In addition, a band below 36 85286, 23877755, 19368702). Plea	normally expressed in cellular cytoplasm o dition of post-translational modifications DH of human origin. It can recognize the 36 kDa can always be detected as the isoform o ase note that some physiological factors, suc pes.	
Notable Publications	also loading control in electrophoresi membrane, but can occasionally tran such as S-nitrosylation. This antibody kDa GAPDH protein in most cells/tiss spliced product of GAPDH (PMID: 238 as hypoxia and diabetes, increase GA Author Put	is and Western blotting. GAPDH is slocate to the nucleus after the ac r is raised against full length GAP uses. In addition, a band below 36 85286, 23877755, 19368702). Plea PDH expression in certain cell typ med ID Journal	normally expressed in cellular cytoplasm of dition of post-translational modifications DH of human origin. It can recognize the 36 kDa can always be detected as the isoform a ase note that some physiological factors, success. Application	
Notable Publications	also loading control in electrophoresi membrane, but can occasionally tran such as S-nitrosylation. This antibody kDa GAPDH protein in most cells/tiss spliced product of GAPDH (PMID: 238 as hypoxia and diabetes, increase GA Author Put Guoqing Sun 361	is and Western blotting. GAPDH is slocate to the nucleus after the ac r is raised against full length GAP ues. In addition, a band below 36 85286, 23877755, 19368702). Plea PDH expression in certain cell typ omed ID Journal 138991 Brain Sci	normally expressed in cellular cytoplasm of dition of post-translational modifications DH of human origin. It can recognize the 36 kDa can always be detected as the isoform ase note that some physiological factors, su pes.	
Notable Publications	also loading control in electrophoresi membrane, but can occasionally tran such as S-nitrosylation. This antibody kDa GAPDH protein in most cells/tiss spliced product of GAPDH (PMID: 238 as hypoxia and diabetes, increase GA Author Put Guoqing Sun 361 Juan Zhao 391	is and Western blotting. GAPDH is slocate to the nucleus after the ac r is raised against full length GAP uses. In addition, a band below 36 85286, 23877755, 19368702). Plea PDH expression in certain cell typ med ID Journal	normally expressed in cellular cytoplasm of dition of post-translational modifications DH of human origin. It can recognize the 36 kDa can always be detected as the isoform of ase note that some physiological factors, success. Application	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data







1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human GAPDH (CL594-60004, Clone:1E6D9) (red), or 0.4 ug Mouse IgG2b Isotype Control (CL594-66360-3, Clone: K1188C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).