For Research Use Only

CoraLite®594-conjugated ESRRB Monoclonal antibody

Catalog Number: CL594-66818



Basic Information

Catalog Number: GenBank Accession Number:

CL594-66818 BC131517 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 2103 Nanodrop: **UNIPROT ID:** 095718 Mouse Full Name:

Isotype: estrogen-related receptor beta

lgG1 Calculated MW: Immunogen Catalog Number: 508 aa, 56 kDa AG18626 Observed MW: 47 kDa

Purification Method: Protein G purification

CloneNo.:

3E10C1

Recommended Dilutions: IF/ICC 1:300-1:1200

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

IF/ICC

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: Caco-2 cells,

Background Information

Estrogen related receptor beta (ESRRB), a NR3 Steroid Receptor, has been shown to affect early placental development. Also it can act as a repressor of transcriptional activity mediated by the glucocorticoid receptor (GR). It binds as a monomer to the extended half-site TNAAGTGCA and as a homodimer to the estrogen response element (ERE), palindromic thyroid hormone response element (TRE(pal)), and SF-1 response element, but not to the glucocorticoid response element (GRE). Estrogen related receptor beta mutations lead to abnormal development of the chorion and defective diploid trophoblast proliferation, and are lethal during midgestation. Expression has been documented in mice in the extra embryonic ectoderm that gives rise to the chorion. ESTs have been isolated from normal human eye, lung, and testis libraries. The ligand for the receptor is PPARgamma coactivator 1 (PGC-1) beta. ESRRB exists various isoforms and molecular weight of each isoform is 48 kDa,55 kDa and 56 kDa.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using Coralite®594 ESRRB antibody (CL594-66818, Clone: 3E10C1) at dilution of 1:600.



1x10^6 Caco-2 cells were intracellularly stained with 0.4 ug Coralite® 594 ESRRB Monoclonal Antibody (CL594-66818, Clone:3E10C1)(red), or 0.4 ug Coralite® 594 Mouse IgG1 Isotype Control (1F8D3) (CL594-66360-1, Clone: 1F8D3) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).