For Research Use Only

## CoraLite® Plus 647-conjugated Nanog Polyclonal antibody

www.ptglab.com

Catalog Number: CL647-14295

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL647-14295 BC160187 GeneID (NCBI):

100ul , Concentration: 1000  $\mu g/ml$  by 79923 Nanodrop:

**UNIPROT ID:** Q9H9S0 Rabbit

Full Name: Isotype: Nanog homeobox IgG Calculated MW: Immunogen Catalog Number: 35 kDa

AG5645 Observed MW:

35-40 kDa

**Purification Method:** 

Antigen affinity purification Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human, mouse, rat

**Background Information** 

Nanog is a member of the homeobox family of DNA binding transcription factors and has been shown to maintain embryonic stem (ES) cell self-renewal independently of leukemia inhibitory factor (LIF)/Stat3. Nanog mRNA is present in pluripotent mouse and human cell lines, and absent from differentiated cells. Functionally, Nanog works together with other key pluripotent factors (Oct4, Sox2, and Lin28) to reprogram human fibroblasts and generate induced pluripotent stem (iPS) cells. These key factors form a regulatory network to support or limit each other's expression level, which maintains the properties of ES cells. Affinity purified rabbit anti-Nanog can be used to demonstrate pluripotency of ES and IPS cells. There are two kinds of variants could recognized by NANOG, one is normal form (~39kd), the other is post-translation modified form (~48kd) (21136380). Nanog exists two isoforms with molecular weight 34.4 kDa and 31.9 kDa. (PMID: 21969378)

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 NCCIT cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human Nanog (CL647-14295) (red), or 0.2 ug isotype control(blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).