

For Research Use Only

# CoraLite® Plus 647 Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody



Catalog Number: **CL647-65559**

## Basic Information

<b>Catalog Number:</b> CL647-65559	<b>GenBank Accession Number:</b> BC025715	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100tests , 5 µl/test	<b>GeneID (NCBI):</b> 925	<b>CloneNo.:</b> UCHT4
<b>Source:</b> Mouse	<b>ENSEMBL Gene ID:</b> ENSG00000153563	<b>Excitation/Emission maxima wavelengths:</b> 654 nm / 674 nm
<b>Isotype:</b> IgG2a	<b>UNIPROT ID:</b> P01732	
	<b>Full Name:</b> CD8a molecule	
	<b>Calculated MW:</b> 235 aa, 26 kDa	

## Applications

**Tested Applications:**  
FC

**Species Specificity:**  
Human

## Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 $\alpha$  or as a heterodimer of CD8 $\alpha$  and CD8 $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 $\alpha\beta$  heterodimer while CD8 $\alpha\alpha$  homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR)  $\gamma\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

## Storage

**Storage:**  
Store at 2-8°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.09% sodium azide and 0.5% BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with FITC Anti-Human CD3 and 5 ul CoraLite® Plus 647 Anti-Human CD8 Mouse Recombinant Antibody (CL647-65559, Clone:UCHL1), or CoraLite® Plus 647 Mouse IgG2a Isotype Control (CL647-65208, Clone: C.1.18.4). Cells were not fixed. Lymphocytes were gated.