

For Research Use Only

# CoraLite® Plus 750-conjugated VDAC1/Porin Monoclonal antibody

Catalog Number:CL750-66345



## Basic Information

<b>Catalog Number:</b> CL750-66345	<b>GenBank Accession Number:</b> NM_003374	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 7416	<b>CloneNo.:</b> 1E2C7
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P21796	<b>Recommended Dilutions:</b> WB 1:1000-1:4000
<b>Isotype:</b> IgG3	<b>Full Name:</b> voltage-dependent anion channel 1	<b>Excitation/Emission maxima wavelengths:</b> 755 nm / 780 nm
	<b>Calculated MW:</b> 31 kDa	
	<b>Observed MW:</b> 35-37 kDa	

## Applications

<b>Tested Applications:</b> WB	<b>Positive Controls:</b> WB : HeLa cells, Jurkat cells, HEK-293 cells
<b>Species Specificity:</b> human, mouse, rat	

## Background Information

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904).


## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-66345 (VDAC1/Porin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.