

For Research Use Only

CoraLite® Plus 750-conjugated FMO1 Recombinant antibody

Catalog Number: CL750-83156-3



Basic Information

Catalog Number:

CL750-83156-3

Size:

100ul , Concentration: 1000 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG34267

GenBank Accession Number:

NM_001282692

GeneID (NCBI):

2326

UNIPROT ID:

Q01740

Full Name:

flavin containing monooxygenase 1

Calculated MW:

60 kDa

Observed MW:

53-60 kDa

Purification Method:

Protein A purification

CloneNo.:

230455B11

Recommended Dilutions:

WB 1:1000-1:4000

Excitation/Emission maxima wavelengths:

755 nm / 780 nm

Applications

Tested Applications:

WB

Species Specificity:

Human, Mouse, Rat

Positive Controls:

WB : mouse liver tissue, human liver tissue

Background Information

Flavin-containing monooxygenase 1 (FMO1) belongs to the FMO family. FMO1 is an enzyme that catalyzes the oxygenation of a series of sulfur-containing and nitrogen-containing substrates, and thus is involved in a series of drugs and xenobiotics metabolism. FMO1 can regulate PPAR- α and mediate metabolic processes. FMO1 is expressed at relatively high levels in human fetal liver and also represents the major FMO enzyme in the adult human and rabbit intestine and kidney (PMID: 11809920, 30757917, 37553313). Human FMO1 has 2 isoforms with the molecular weights of 53 and 60 kDa. The observed molecular weight of FMO1 is 53 kDa, which might be the unmodified form of FMO1 (PMID: 9778310).

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-83156-3 (FMO1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.