For Research Use Only

FITC Plus Anti-Human CD18 (TS1/18) Mouse IgG2a Recombinant Antibody

Catalog Number:FITC-65595



Basic Information

Catalog Number:

BC005861

Purification Method: Protein A purification

Size:

GeneID (NCBI): 3689 CloneNo.:

100tests . 5 ul/test

FITC-65595

3009

TS1/18

495 nm / 524 nm

Source: Mouse ENSEMBL Gene ID: ENSG00000160255

Excitation/Emission maxima wavelengths:

Isotype:

Full Name:

lgG2a

integrin, beta 2 (complement

GenBank Accession Number:

component 3 receptor 3 and 4 subunit)

Calculated MW:

85 kDa

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD18, also known as integrin beta-2, is a transmembrane protein that forms heterodimers with CD11a, CD11b, CD11c, CD11d (PMID: 1968349). CD11a/CD18 (LFA-1) is expressed by all leukocytes, while CD11b/CD18 (Mac-1) and CD11c/CD18 (p150,95) are normally restricted to expression on monocytes, macrophages, polymorphonuclear lymphocytes (PMNs), and natural killer cells (PMID: 1968349; 3109455; 10946284). CD18 plays an important role in mediating cell adhesion during immune response and defects of CD18 cause leukocyte adhesion deficiency (PMID: 3594570).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with APC Anti-Human CD45 (HI30) (APC-65109, Clone: HI30) and 5 ul FITC Plus Anti-Human CD18 (TS1/18) Mouse IgG2a Recombinant Antibody (FITC-65595, Clone:TS1/18), or FITC Plus Mouse IgG2a Isotype Control (C1.18.4) (FITC-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD18 (T51/18) Mouse IgG2a Recombinant Antibody (FITC-65595, Clone:TS1/18)(red), or FITC Plus Mouse IgG2a Isotype Control (C1.18.4) (FITC-65208, Clone: C1.18.4)(blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.