For Research Use Only

FITC Plus Anti-Human CD81 Rabbit Recombinant Antibody

Catalog Number:FITC-98202



Basic Information

Catalog Number: GenBank Accession Number: FITC-98202 NM_004356.4

GeneID (NCBI): Size:

100tests, 5 ul/test

ENSEMBL Gene ID: Source: Rabbit ENSG00000110651

UNIPROT ID: Isotype: P60033

Full Name: Immunogen Catalog Number: EG1036

CD81 molecule Calculated MW:

26 kDa

Purification Method:

Protein A purification

CloneNo.: 241722G4

Recommended Dilutions:

FC: 5 ul per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 495 nm / 524 nm

Applications

Tested Applications:

Species Specificity:

human

Positive Controls:

FC: human PBMCs,

Background Information

CD81 (also known as TAPA1 or TSPAN28) is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Many of these members are expressed on leukocytes and have been implicated in signal transduction, cell-cell interactions, and cellular activation and development. CD81 is involved in signal transduction and cell adhesion in the immune system (PMID: 9597125). CD81 has also been identified as en essential receptor for HCV (hepatitis C virus) (PMID: 21428934).

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD81 Rabbit RecAb (FITC-98202, Clone: 241722G4) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD81 Rabbit RecAb (FITC-98202, Clone: 241722G4) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.