

For Research Use Only

FITC Plus Anti-Mouse Ki-67 Rabbit Recombinant Antibody

Catalog Number: FITC-98224



Basic Information

Catalog Number:

FITC-98224

Size:

100ug , 500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG1548

GenBank Accession Number:

NM_001081117.2

GeneID (NCBI):

17345

UNIPROT ID:

E9PVX6-1

Full Name:

antigen identified by monoclonal antibody Ki 67

Calculated MW:

351kDa

Purification Method:

Protein A purification

CloneNo.:

241768E2

Recommended Dilutions:

FC (Intra): 0.25 ug per 10⁶ cells in 100 µl suspension

Excitation/Emission maxima wavelengths:

495 nm / 524 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

mouse

Positive Controls:

FC (Intra) : Con A treated C57BL/6 mouse splenocytes,

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki-67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse Ki-67 Rabbit RecAb (FITC-98224, Clone: 241768E2). Cells were co-stained with APC Anti-Mouse CD3 (17A2) (APC-65077, Clone: 17A2). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

1x10⁶ Con A treated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse Ki-67 Rabbit RecAb (FITC-98224, Clone: 241768E2) (red) or FITC Plus Rabbit IgG Isotype Control RecAb (FITC-98136, Clone: 240953C9) (blue). 1x10⁶ untreated C57BL/6 mouse splenocytes were intracellularly stained with 0.25 ug FITC Plus Anti-Mouse Ki-67 Rabbit RecAb (FITC-98224, Clone:241768E2) (black, dashed). Cells were fixed and

