PE-conjugated PLOD2-Specific Polyclonal antibody

Catalog Number: PE-21214

Featured Product

Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: PE-21214	GenBank Accession Number: BC037169	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 200 µg/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG15614	GeneID (NCBI): 53552 UNIPROT ID: 000469 Full Name: procollagen-lysine, 2-oxoglutarate 5 dioxygenase 2 Calculated MW: 758 aa, 85 kDa Observed MW: 87 kDa	Excitation/Emission maxima wavelengths: 496 nm, 565 nm / 578 nm
Applications	Tested Applications: FC (Intra) Species Specificity: human, mouse, rat		
Background Information	PLOD2, also named as LH2, forms hydroxylysine residues in -Xaa-Lys-Gly- sequences in collagens. It is a potential novel prognostic factor for HCC patients following surgery. Among the PLOD genes, PLOD2 contributes to cancer prognosis and angiogenesis. Several authors have reported that PLOD2 expression might provide prognostic information about malignant tumours such as glioblastoma. PLOD2 expression is a useful biomarker for the effects of antiangiogenic treatment for malignancy. (PMID:22098155). It has 2 isoforms produced by alternative splicing and seven glycosylation sites. This antibody is specific to PLOD2.		


Storage

Storage: Store at -20°C. Avoid exposure to light. Storage Buffer: PBS with 0.1% sodium azide and 0.5% BSA. Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.4 ug PE-conjugated PLOD2-Specific Polyclonal antibody (PE-21214)(red), or 0.4 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).