

For Research Use Only.
Not For Use In Diagnostics.

HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody

Catalog Number:RGAM701



www.ptglab.com

Information

Catalog Number:

RGAM701

Host:

Goat

Applications:

ELISA, WB, Dot blot

Reactivity:

Mouse

Physical State:

Liquid

Conjugation:

HRP

Clonality:

Recombinant monoclonal

Recommended Dilutions

1:10,000-1:100,000 for ELISA

1:10,000-1:200,000 for western blotting with ECL substrates (1:10,000-1:40,000 is suggested for most systems).

Safety Notes

This product is for research use only, not for diagnostic or therapeutic use.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 50% glycerol, 10 mg/mL BSA, 0.1% Proclin-300, pH 7.4.

Aliquoting is unnecessary for -20°C storage

Purity

The antibody was purified from culture media supernatant by immunoaffinity chromatography using Protein G beads.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB
(1-888-478-4522)
(toll free in USA),
or 1(312) 455-8498
(outside USA)


E : proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS-PAGE followed by western blot with Actin mouse monoclonal antibody (60008-1-Ig, IgM). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. Note that a higher signal is obtained using RGAM001 as it recognizes multiple epitopes.




Various lysates were subjected to SDS-PAGE, followed by western blot with Beta-Tubulin mouse monoclonal antibody (66240-1-Ig, IgG2a). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgG2a.




Various lysates were subjected to SDS-PAGE followed by western blot with FAK mouse monoclonal antibody (66258-1-Ig, IgG2c). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgG2c.




Various lysates were subjected to SDS-PAGE followed by western blot with Gelsolin mouse monoclonal antibody (66280-1-Ig, IgA). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgA.




Various lysates were subjected to SDS-PAGE, followed by western blot with NRF 1 mouse monoclonal antibody (66832-1-Ig, IgG2b). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgG2b.



Various lysates were subjected to SDS-PAGE followed by western blot with EIF 3E mouse monoclonal antibody (67095-1-Ig, IgG1). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgG1.



Various lysates were subjected to SDS-PAGE followed by western blot with ZNF 174 mouse monoclonal antibody (68426-1-Ig, IgG3). HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701) and Multi-rAb HRP-Goat Anti-Mouse IgG (H+L) Recombinant Secondary Antibody (RGAM001) were used at 1: 20000 for detection. The result shows that RGAM701 does not react with mouse IgG3.



Direct ELISA was performed by coating mouse monoclonal antibodies with different isotypes followed by signal development using HRP-Goat Anti-Mouse IgM-Specific Recombinant Secondary Antibody (RGAM701). The result indicates that RGAM701 strongly binds to Mouse IgM and does not react with Mouse IgGs or mouse IgA.