For Research Use Only

Multi-rAb™ TSG101 Multi-Recombinant antibody

www.ptglab.com

Purification Method:

WB 1:5000-1:50000

protein lysate

IHC 1:300-1:1200

IF/ICC 1:200-1:800

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Catalog Number: RMX00026

Basic Information

Applications

Catalog Number: GenBank Accession Number:

RMX00026 BC002487 GeneID (NCBI): 100ul , Concentration: 500 $\mu g/ml$ by

Nanodrop: **UNIPROT ID:** Source: Q99816 Rabbit Full Name:

Isotype: tumor susceptibility gene 101 IgG

Calculated MW: 44 kDa Observed MW:

44 kDa

Positive Controls:

WB, IHC, IF/ICC, IP, ELISA WB: HeLa cells, Jurkat cells, K-562 cells, NIH/3T3 cells,

> HSC-T6 cells IP: HeLa cells,

Note-IHC: suggested antigen retrieval with IHC: human colon cancer tissue, TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate IF/ICC: HeLa cells, buffer pH 6.0

Background Information

TSG101(Tumor susceptibility gene 101 protein) is essential for endosomal sorting, membrane receptor degradation and the final stages of cytokinesis. It plays a crucial role for cell proliferation and cell survival. TSG101 has been identified as a candidate tumor suppressor gene and belongs to the ubiquitin-conjugating enzyme family. TSG101 is a marker for exosome. This protein has 2 isoforms produced by alternative splicing with the molecular mass of 44 and 32 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

Tested Applications:


Species Specificity:

human, mouse, rat


PBS with 0.02% sodium azide and 50% glycerol. Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with RMX00026 (TSG101 antibody) at dilution of 1:12000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using RMX00026 (TSG101 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using TSG101 antibody (RMX00026) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



IP result of anti-TSG101 (IP:RMX00026, 4ug; Detection:RMX00026 1:10000) with HeLa cells lysate 1480 ug.