

iST V5-Trap® IP-MS Sample Preparation Kit


Product Code: v5tak-iST



Introduction

The PreOmics®/ChromoTek® iST V5-Trap® Kit enables efficient and fast immunoprecipitation of V5-tagged proteins and their interaction factors, as well as the subsequent sample preparation for mass spectrometry-based proteomics. In the first step, the kit's V5-Trap® utilizes small recombinant nanobodies/VHHs covalently coupled to agarose beads for pull-down. Then in a second step, this kit provides buffers, cartridges and enzymes to denature, reduce, alkylate, and digest V5-tagged proteins and their interaction factors, and to perform a final peptide cleanup step in a streamlined manner. For sample-specific protocols and optimization contact us or visit our websites www.ptglab.com or www.preomics.com.

Procedure



Kit Contents

Component	Cap	Quantity	Buffer Properties				Description	Storage
			Organic	Acidic	Basic	Volatile		
V5-Trap Agarose	green/white	0.25 ml slurry					Anti-V5-tag nanobody coupled to agarose beads (v5ta-10; i.e. 8 reactions plus 2 controls).	4°C
DIGEST	red	2x					Enzyme <i>Trypsin</i> -mix to digest proteins.	-20°C
RESUSPEND	yellow	1x 1 ml				✓	Protease reconstitution buffer for enzymes.	RT
LYSE	brown	1x 1 ml			✓		Denature, reduce and alkylate proteins.	RT
STOP	black	1x 1 ml	✓	✓		✓	Stop the enzymatic activity.	RT
WASH 1	blue	1x 2 ml	✓	✓		✓	Clean up peptides from hydrophobic contaminants.	RT
WASH 2	green	1x 2 ml		✓		✓	Clean up peptides from hydrophilic contaminants.	RT
ELUTE	violet	1x 2 ml	✓		✓	✓	Elute the peptides from the cartridge.	RT
LC-LOAD	white	1x 1 ml		✓		✓	Load peptides on reversed-phase LC-MS column.	RT
CARTRIDGE		8x					Cartridge for 1 to 100 µg protein starting material.	RT
WASTE		8x					Tube for collecting waste after washing steps.	RT
COLLECTION		8x					Tube for collecting peptides after elution.	RT
ADAPTER		8x					Enables placing a cartridge into a tube.	RT

The iST V5-Trap® Kit contains sufficient V5-Trap® Agarose beads for up to 10 immunoprecipitations. Out of these 10 reactions, two shall be used for negative and positive controls by immunoblot analysis. The iST V5-Trap® Kit can be used for up to 8 iST proteomic sample preparation reactions.

iST V5-Trap® IP-MS Sample Preparation Kit

Product Code: v5tak-iST



V5-Trap® Agarose Bead Properties

Bead size: ~ 90 µm (cross-linked 4% agarose beads)

Storage buffer: 20% EtOH

Binding capacity: 25 µL V5-Trap Agarose slurry binds 15 µg of recombinant V5-tagged protein

V5-Trap® Agarose Stability and Storage

Shipped at ambient temperature. Upon receipt store at +4°C. Stable for 1 year. Do not freeze.

Required Equipment

Common lab equipment is required for the sample preparation:

EQUIPMENT	QUANTITY AND DESCRIPTION
Pipette	Careful sample handling and pipetting reduces contaminations and improves quantification.
Sample	Lyophilized protein or pelleted cells. For other materials ask us for adapted protocols.
Heating Block	Heating shakers are recommended to help protein denaturation and during digestion.
Centrifuge	Microcentrifuge for 1.5 ml reaction tubes is necessary for loading, washing and elution.
Speed-Vac	Vacuum manifolds evaporate volatile buffers from the eluate before LC-MS.
Ultrasonic Bath	Optional: can be used to resuspend peptides.
Reaction Tubes	1.5 mL plasticware reaction tubes, e.g. from Eppendorf. We recommend low-binding protein tubes.
Ice Bucket	Keep required solutions and protein samples on ice.

Required Solutions (Not Included)

Suggested buffer compositions for mammalian cells. For other cell types like yeast, plants, drosophila, etc. please use equivalent cell lysis buffer. We advise to use MS-grade chemicals to prepare all buffers.

IP LYSIS BUFFER	Selection of IP LYSIS Buffer: For <u>cytoplasmic proteins</u>, re-suspend cell pellet in 200 µl ice-cold IP LYSIS buffer (not included) by pipetting up and down. Supplement IP LYSIS buffer with protease inhibitor cocktail and 1 mM PMSF. Composition: 10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.5 mM EDTA; 0.5% NP-40 For <u>nuclear and cytoplasmic proteins</u>, resuspend cell pellet in 200 µl ice-cold RIPA buffer (not included) supplemented with DNaseI (f.c. 75-150 U/ml), MgCl₂ (f.c. 2.5 mM), protease inhibitor cocktail and PMSF (f.c. 1 mM). Composition: 10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.5 mM EDTA; 0.1 % SDS; 1 % Triton X-100; 1 % Deoxycholate
IP WASH I	10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0.25% NP-40
IP WASH II	10 mM Tris/Cl pH 7.5; 150 mM NaCl
PBS	Phosphate-buffered saline: 137 mM NaCl; 2.7 mM KCl; 10 mM Na ₂ HPO ₄ ; 1.8 mM KH ₂ PO ₄ ; adjust to a final pH of 7.4

Immunoprecipitation using V5-Trap® Agarose

Keep solutions and protein samples on ice.

1. Harvest

- 1.1 Use of $\sim 10^6$ - 10^7 mammalian cells (approx. one 10-cm dish) expressing the V5-tagged protein of interest for one immunoprecipitation reaction. Aspirate growth medium, add 1 mL ice-cold PBS to cells and scrape cells from dish to harvest adherent cells.
- 1.2 Transfer cells to a pre-cooled 1.5 mL tube, spin at 500 rcf for 3 min at 4°C and discard supernatant.
- 1.3 Wash cell pellet twice with ice-cold **PBS**, gently resuspending the cells.

2. IP Lysis

- 2.1 Resuspend cell pellet in 200 μ L ice-cold IP LYSIS buffer supplemented with protease inhibitors and 1 mM PMSF by pipetting or using a syringe. See IP LYSIS buffer composition on page 2.
- 2.2 Place the tube on ice for 30 min with extensively pipetting every 10 min.
- 2.3 Centrifuge cell lysate at 20,000 rcf for 10 min at 4°C.
- 2.4 Transfer supernatant to a pre-cooled 1.5 mL tube. Add 300 μ L IP WASH II buffer supplemented with protease inhibitors and 1 mM PMSF to lysate.
- 2.5 For negative/positive control save 50 μ L diluted lysate for immunoblot analysis (input).

3. Equilibrate Beads

- 3.1 Resuspend **V5-Trap Agarose beads**  by pipetting and transfer 25 μ L bead slurry into 500 μ L ice-cold **IP WASH I buffer**.
- 3.2 Resuspend beads and pipette 25 μ L bead slurry into 500 μ L ice-cold **IP WASH I buffer**.
- 3.3 Centrifuge at 2,500 rcf for 2 min at +4°C. Discard supernatant and repeat wash once.

4. Bind Proteins

- 4.1. Add diluted lysate (step 2.4) to equilibrated **V5-Trap Agarose Beads** (step 3.3).
- 4.2. Tumble end-over-end for 1 hour at 4°C.
- 4.3. Centrifuge at 2,500 rcf for 2 min at +4°C. If required, save 50 μ L supernatant for immunoblot analysis. Discard remaining supernatant.

5. Wash Beads

- 5.1. Resuspend **V5-Trap Agarose beads** in 500 μ L ice-cold **IP WASH I buffer**. Centrifuge at 2,500 rcf for 2 min at +4°C.
- 5.2. Discard supernatant and repeat this wash step once.
Optional: Increase salt concentration up to 500 mM for higher stringency washing.
- 5.3. Resuspend beads in 500 μ L ice-cold **IP WASH II buffer**. Centrifuge at 2,500 rcf for 2 min at +4°C.
- 5.4. Discard supernatant, repeat this wash step three times. Transfer beads to a **fresh tube** at last wash step.
Optional: Increase salt concentration up to 500 mM for higher stringency washing. Do not include detergents.
- 5.5. For negative/positive control (bound protein) resuspend 25 μ L beads in 100 μ L 2x SDS-sample buffer. Boil resuspended beads for 10 min at 95°C to dissociate immunocomplexes from V5-Trap Agarose beads. Beads can be collected by centrifugation at 2,500 rcf for 2 min at 4°C and SDS-PAGE is performed with the supernatant.
- 5.6. Discard supernatant and directly continue with the proteomic sample preparation using the iST part of kit as outlined on next page.

iST V5-Trap® IP-MS Sample Preparation Kit

Product Code: v5tak-iST



Proteomic sample preparation using iST

6. Denature Proteins

- 6.1. Add 50 µL **LYSE** ● to the washed **beads** and place it in a pre-heated HEATING BLOCK (60 °C; 1,000 rpm; 10 min).
- 6.2. Optional: Spin down droplets (RT; max. 300 rcf; 10 sec).

7. Digest

- 7.1. Add 210 µL **RESUSPEND** ● to **DIGEST** ● (1 tube for 4 reactions), shake (RT; 500 rpm; 10 min), pipette up/down.
 - 7.2. Add 50 µL **DIGEST** ● to **beads** and place it in a pre-heated HEATING BLOCK (37 °C; 500 rpm; 1-3 hours).
 - 7.3. Add 100 µL **STOP** ● to **beads** (precipitation may happen), shake (RT; 500 rpm; 1 min), pipette up/down.
- *SP*

8. Purify

- 8.1. Use **ADAPTER** to place **CARTRIDGE** in **WASTE** tube. Label all tubes.
- 8.2. Centrifuge **beads** (RT; 2,500 rcf; 2 min).
- 8.3. Transfer the complete supernatant (combined LYSE/DIGEST/STOP buffers) to **CARTRIDGE**. Discard beads.
- 8.4. Spin **CARTRIDGE** in CENTRIFUGE (RT; 3,800 rcf; 1-3 min). If needed, adjust values to ensure complete flow-through.
- 8.5. Add 200 µL **WASH 1** ● to **CARTRIDGE**, repeat step 8.4., discard flow-through.
- 8.6. Add 200 µL **WASH 2** ● to **CARTRIDGE**, repeat step 8.4., discard flow-through. *SP*
- 8.7. Use **ADAPTER** to place **CARTRIDGE** in a fresh **COLLECTION** tube. Label all tubes.
- 8.8. Add 100 µL **ELUTE** ● to **CARTRIDGE**, repeat step 8.3., keep flow-through in **COLLECTION** tube.
- 8.9. Repeat step 8.8., keep flow-through in the same **COLLECTION** tube.
- 8.10. Remove the **CARTRIDGE** and place **COLLECTION** tube in a SPEED-VAC (45 °C; until completely dry).
- 8.11. Add **LC-LOAD** ○ to **COLLECTION** tube. Aim for 1 g/L concentration (e.g. 50 µL to 50 µg pulled-down protein).
- 8.12. Sonicate **COLLECTION** tube in an ULTRASONIC BATH (5 min) or shake (RT; 500 rpm; 5 min). *SP*

***SP* - Storage Point:** At this point, close the tube or **CARTRIDGE**. Lysates and peptides can be frozen at -20 °C. Storage of peptides should not exceed 2 weeks at -20 °C. For extended storage, finish the protocol and store at -80 °C.

Data Analysis

Consider the following as fixed modifications in your database search:

Modification	Description	Composition	Specificity	Mass	Unimod #
ALKYLATION	Carbamidomethyl on Cysteine	C ₂ H ₃ NO	[C]	+57 Da	4

iST V5-Trap® IP-MS Sample Preparation Kit

Product Code: v5tak-iST



Contact

Proteintech North America (HQ)

Proteintech Group, Inc.
5500 Pearl Street, Suite 400
Rosemont, IL 60018 USA

1-888-472-4522

proteintech@ptglab.com

Proteintech Europe

Transmission (6th Fl)
6 Atherton Street
M3 3GS, Manchester, UK

+44 161 839 3007

europe@ptglab.com

ChromoTek & Proteintech Germany

Fraunhoferstr. 1
82152, Planegg-Martinsried
Germany

+49 89 124 148 850

germany@ptglab.com

Disclaimer

For research use only (RUO), not for diagnostic or therapeutic use.

ChromoTek® and V5-Trap® are registered trademarks of ChromoTek GmbH, part of Proteintech Group.

PreOmics® is a registered trademark of PreOmics GmbH in the US, EU, and UK.